

La tabla incluye todos los taxones de aves playeras de ocurrencia regular en las Américas (i.e. excluye aquellas especies consideradas errantes en la región)

Especie	Población biogeográfica	Estimación	Punto medios <sup>1</sup>	Fuente	Nivel umbral del 1%
<i>Jacana spinosa</i>	<i>spinosa</i>	48,000 - 170,000	109,000	WPE5	1000
	<i>gymnostoma</i>	NA			
	<i>violacea</i>	<10,000		WPE5	100
<i>Jacana jacana</i>	<i>jacana</i>	100,000 – 1,000,000		WPE5	10,000
	<i>hypomelaena</i>	NA			
	<i>melanopygia</i>	NA			
	<i>intermedia</i>	NA			
	<i>scapularis</i>	NA			
	<i>peruviana</i>	NA			
<i>Rostratula semicollaris</i>		NA			
<i>Haematopus leucopodus</i>	S South America	25,000 - 100,000		WPE5	630
	Falkland Is. (Malvinas)	21,000 - 39,000	30,000	WPE5	300
<i>Haematopus ater</i>	S South America	10,000-100,000		WPE5	500
	Falkland Is. (Malvinas)	12,000 - 24,000	18,000	WPE5	180
<i>Haematopus bachmani</i>		10,000		Andres <i>et al.</i> 2012	100
<i>Haematopus palliatus</i>	<i>palliatus</i> (Total)	20,266		Clay <i>et al.</i> 2009	200
	<i>palliatus</i> - USA	11,284		Brown <i>et al.</i> 2005, Schulte unpub. data	110
	<i>palliatus</i> – Caribbean/ N S. America	750		Clay <i>et al.</i> 2014	8
	<i>palliatus</i> – E. South America	4,910		Clay <i>et al.</i> 2014	50
	<i>frazari</i>	3,000		Clay <i>et al.</i> 2009	30
	<i>pitanay</i>	10,000-15,000	12,500	Clay <i>et al.</i> 2009	130
	<i>durnfordi</i>	10,000-15,000	12,500	Clay <i>et al.</i> 2009	130
	<i>galapagensis</i>	300		Clay <i>et al.</i> 2009	3
<i>Himantopus mexicanus</i>	<i>mexicanus</i>	175,000-1,000,000		WPE5	6,000
	<i>knudseni</i>	2,100		Andres <i>et al.</i> 2012	20
	<i>melanurus</i>	100,000-1,000,000		WPE5	5,000
<i>Recurvirostra americana</i>		450,000		Andres <i>et al.</i> 2012	4,500
<i>Recurvirostra andina</i>		<10,000		Boyla & Estrada 2005	50

<sup>1</sup> Presentados son los rangos dados en Andres *et al.* 2012 y otros rangos relativamente precisos. Los puntos medios para los rangos proporcionados por WPE no se consideran aquí.

<i>Burhinus bistrriatus</i>	<i>bistrriatus</i>	2,600 - 5,450	4,025	WPE5	40
	<i>pediacus</i>	10,000		WPE5	100
	<i>vocifer</i>	10,000		WPE5	100
	<i>dominicensis</i>	NA			
<i>Burhinus superciliaris</i>		<10,000		Boyla & Estrada 2005	50
<i>Vanellus cayanus</i>		25,000 – 100,000		WPE5	630
<i>Vanellus chilensis</i>	<i>chilensis</i>	>1,000,000		WPE5	20,000
	<i>fretensis</i>	>1,000,000		WPE5	20,000
	<i>cayennensis</i>	25,000 – 100,000		WPE5	630
	<i>lampronotus</i>	NA			
<i>Vanellus resplendens</i>		10,000 - 25,000		WPE5	150
<i>Pluvialis fulva</i>	Alaska (Br)	35,000-50,000	42,500	Andres <i>et al.</i> 2012	425
<i>Pluvialis dominica</i>		294,200-705,800	500,000	Andres <i>et al.</i> 2012	5,000
<i>Pluvialis squatarola</i>	<i>squatarola</i> (Alaska; Br)	134,000-391,500	262,700	Andres <i>et al.</i> 2012	2600
	<i>cynosurae</i>	100,000		Andres <i>et al.</i> 2012	1,000
<i>Charadrius hiaticula</i>	<i>psammodroma</i>	2,000		Andres <i>et al.</i> 2012	20
<i>Charadrius semipalmatus</i>		200,000		Andres <i>et al.</i> 2012	2,000
<i>Charadrius wilsonia</i>	<i>wilsonia</i> (Total)	13,550 - 14,650	14,100	Zdravkovic 2013	140
	<i>wilsonia</i> - USA	8,000 – 8,600	8,300	Zdravkovic 2013	85
	<i>wilsonia</i> - Caribbean	3,600 – 4,000	3,800	Zdravkovic 2013	40
	<i>cinnamoniis</i>	6,500 - 8,500	7,500	Zdravkovic 2013	75
	<i>beldingi</i>	6,500 - 8,500	7,500	Zdravkovic 2013	75
	<i>crassirostris (brasiliensis)</i>	NA			
<i>Charadrius vociferus</i>	<i>vociferus</i>	2,000,000		Andres <i>et al.</i> 2012	20,000
	<i>ternominatus</i>	<10,000		WPE5	50
	<i>peruvianus</i>	10,000-25,000		Boyla & Estrada 2005	150
<i>Charadrius melodus</i>	<i>melodus</i> (Atlantic)	3,600		Andres <i>et al.</i> 2012	35
	<i>circumcinctus</i> (Great Lakes; Br)	108		Andres <i>et al.</i> 2012	1
	<i>circumcinctus</i> (Great Plains)	4,700		Andres <i>et al.</i> 2012	45
<i>Charadrius nivosus</i>	<i>nivosus</i> (Interior Atlantic)	16,700-29,200	22,900	Andres <i>et al.</i> 2012	230
	<i>nivosus</i> (Pacific)	2,900		Andres <i>et al.</i> 2012	30
	<i>tenuirostris</i>	<200		Andres <i>et al.</i> 2012	2
	<i>occidentalis</i>	<10,000		WPE5	50
<i>Charadrius collaris</i>	<i>collaris</i>	<10,000		WPE5	50
	<i>gracilis</i>	10,000 - 25,000		Boyla & Estrada 2005	150
<i>Charadrius alticola</i>		25,000 – 1,000,000		WPE5	10,000
<i>Charadrius falklandicus</i>	S South America	25,000 – 100,000		WPE5	630
	Falkland Is. (Malvinas)	21,000 - 39,000	30,000	WPE5	300
<i>Charadrius montanus</i>		20,000		Andres <i>et al.</i> 2012	200
<i>Charadrius modestus</i>	S South America	100,000-1,000,000		WPE5	5,000
	Falkland Is. (Malvinas)	33,000 - 63,000	48,000	WPE5	480
<i>Oreopholus ruficollis</i>	<i>ruficollis</i>	<10,000		WPE5	50
	<i>pallidus</i>	<10,000		Boyla & Estrada 2005	50
<i>Phegornis mitchellii</i>		<10,000		WPE5	50

<i>Pluvianellus socialis</i>		1,500		WPE5	15
<i>Chionis albus</i>		10,000 - 25,000		Lesterhuis & Clay 2012	150
<i>Scolopax minor</i>	Atlantic North America (Br)	3,000,000-4,000,000	3,500,000	Andres <i>et al.</i> 2012	35,000
	Inland E North America (Br)				
<i>Gallinago delicata</i>		2,000,000		Andres <i>et al.</i> 2012	20,000
<i>Gallinago paraguaiiae</i>	<i>paraguaiiae</i>	25,000-100,000		WPE5	630
	<i>magellanica</i> (S South America)	25,000 – 1,000,000		WPE5	10,000
	<i>magellanica</i> (Falkland Is. (Malvinas))	15,000 - 27,000	21,000	WPE5	210
<i>Gallinago andina</i>		10,000 – 25,000		Boyla & Estrada 2005	150
<i>Gallinago nobilis</i>		<10,000		Cisneros-Heredia 2006	50
<i>Gallinago undulata</i>	<i>undulata</i>	<10,000		Boyla & Estrada 2005	50
	<i>gigantea</i>	10,000 – 25,000		Boyla & Estrada 2005	150
<i>Gallinago stricklandii</i>		<10,000		WPE5	50
<i>Gallinago jamesoni</i>		3,000 – 10,000	6,500	WPE5	65
<i>Gallinago imperialis</i>		<10,000		Boyla & Estrada 2005	50
<i>Limnodromus griseus</i>	<i>griseus</i>	78,000		Andres <i>et al.</i> 2012	1,500
	<i>hendersoni</i>				
	<i>caurinus</i>	75,000			
<i>Limnodromus scolopaceus</i>		500,000		Andres <i>et al.</i> 2012	5,000
<i>Limosa haemastica</i>	Hudson Bay (Br)	56,000		Andres <i>et al.</i> 2012	770
	Alaska (Br)	21,000			
<i>Limosa lapponica</i>	<i>baueri</i>	90,000 (80,000-120,000)		Andres <i>et al.</i> 2012	900
<i>Limosa fedoa</i>	<i>fedoa</i> (SC Canada & NC USA; Br)	170,000		Andres <i>et al.</i> 2012	1,700
	<i>fedoa</i> (James Bay; Br)	2,000			
	<i>beringiae</i>	2,000 (2,000-3,000)			
<i>Numenius borealis</i>		<50		Andres <i>et al.</i> 2012	1
<i>Numenius phaeopus</i>	<i>rufiventris</i>	40,000		Andres <i>et al.</i> 2012	800
	<i>hudsonicus</i>	40,000			
<i>Numenius tahitiensis</i>		10,000		Andres <i>et al.</i> 2012	100
<i>Numenius americanus</i>	<i>americanus</i>	98,000-198,000	140,000	Andres <i>et al.</i> 2012	1,400
<i>Bartramia longicauda</i>		750,000		Andres <i>et al.</i> 2012	7,500
<i>Tringa melanoleuca</i>		137,000		Andres <i>et al.</i> 2012	1,300
<i>Tringa flavipes</i>		660,000		Andres <i>et al.</i> 2012	6,600
<i>Tringa solitaria</i>	<i>solitaria</i>	126,000		Andres <i>et al.</i> 2012	1900
	<i>cinnamomea</i>	63,000			
<i>Actitis macularius</i>		660,000		Andres <i>et al.</i> 2012	6,600
<i>Tringa incana</i>	N North America (Br)	10,000-25,000	17,500	Andres <i>et al.</i> 2012	175
<i>Tringa semipalmatus</i>	<i>semipalmatus</i>	90,000		Andres <i>et al.</i> 2012	2,500
	<i>inornatus</i>	160,000			
<i>Arenaria interpres</i>	<i>interpres</i> (NE Canada; Br)	45,000		Andres <i>et al.</i> 2012	2,500
	<i>interpres</i> (Alaska; Br)	20,000			
	<i>morinella</i>	180,000			
<i>Arenaria melanocephala</i>		76,000-114,000	95,000	Andres <i>et al.</i> 2012	950

<i>Calidris virgata</i>		70,000		Andres <i>et al.</i> 2012	700
<i>Calidris canutus</i>	<i>islandica</i>	80,000		Andres <i>et al.</i> 2012	800
	<i>rufa</i> (Total)	42,000		Andres <i>et al.</i> 2012	420
	<i>rufa</i> - Tierra del Fuego	13,500		R. Morrison in litt. to BirdLife 2015	135
	<i>rufa</i> - N. Brazil	15,500		R. Morrison in litt. to BirdLife 2015	155
	<i>rufa</i> - SE USA	11,000		Calculated from Andres <i>et al.</i> 2012	110
	<i>roselaari</i>	21770		Lyons <i>et al.</i> 2015	210
<i>Calidris alba</i>	N & S America (N-Br)	300,000		Andres <i>et al.</i> 2012	3,000
<i>Calidris pusilla</i>		1,728,400-2,791,600	2,260,000	Andres <i>et al.</i> 2012	22,600
<i>Calidris mauri</i>		3,500,000		Andres <i>et al.</i> 2012	35,000
<i>Calidris minutilla</i>		700,000		Andres <i>et al.</i> 2012	7,000
<i>Calidris fuscicollis</i>		560,100-3,827,900	1,694,000	Andres <i>et al.</i> 2012	17,000
<i>Calidris bairdii</i>		300,000		Andres <i>et al.</i> 2012	3,000
<i>Calidris melanotos</i>		1,129,600-2,070,400	1,600,000	Andres <i>et al.</i> 2012	16,000
<i>Calidris acuminata</i>		16,000-32,000	22,000	Andres <i>et al.</i> 2012	220
<i>Calidris himantopus</i>		418,800-2,068,600	1,243,700	Andres <i>et al.</i> 2012	12,400
<i>Calidris maritima</i>	<i>maritima</i>	10,000		Andres <i>et al.</i> 2012	250
	<i>belcheri</i>	15,000			
<i>Calidris ptilocnemis</i>	<i>ptilocnemis</i>	17,900-21,900	19,800	Andres <i>et al.</i> 2012	1,450
	<i>tschuktschorum</i>	50,000			
	<i>couesi</i>	75,000			
<i>Calidris alpina</i>	<i>arctica</i>	304,000-696,000	500,000	Andres <i>et al.</i> 2012	5,000
	<i>pacifica</i>	550,000		Andres <i>et al.</i> 2012	5,500
	<i>hudsonia</i>	220,700-679,300	450,000	Andres <i>et al.</i> 2012	4,500
<i>Calidris subruficollis</i>		35,000-78,000	56,000	Andres <i>et al.</i> 2012	560
<i>Phalaropus tricolor</i>		1,500,000		Andres <i>et al.</i> 2012	15,000
<i>Phalaropus lobatus</i>	North America (Br)	2,500,000		Andres <i>et al.</i> 2012	25,000
<i>Phalaropus fulicarius</i>	Alaska & Canada (Br)	1,143,700-2,096,300	1,620,000	Andres <i>et al.</i> 2012	16,200
<i>Attagis gayi</i>	<i>gayi</i>	25,000 – 1,000,000		WPES	10,000
	<i>latreilli</i>	< 600		WPES	3
	<i>simonsi</i>	25,000 – 1,000,000		WPES	10,000
<i>Attagis malouinis</i>		10,000 – 25,000		Boyla & Estrada 2005	150
<i>Thinocorus orbignyianus</i>	<i>orbignyianus</i>	10,000 - 25,000		WPES	150
	<i>ingae</i>	10,000 - 25,000		Boyla & Estrada 2005	150
<i>Thinocorus rumicivorus</i>	<i>rumicivorus</i>	100,000-1,000,000		WPES	5,000
	<i>cuneicauda</i>	< 25,000		WPES	125
	<i>bolivianus</i>	10,000 - 25,000		Boyla & Estrada 2005	150
	<i>pallidus</i>	< 10,000		Boyla & Estrada 2005	50

## **Referencias**

- Andres, B.A., Smith, P.A., Morrison, R.I.G., Gratto-Trevor, C.L., Brown, S.C. & Friis, C.A. 2012. Population estimates of North American shorebirds, 2012. *Wader Study Group Bull.* 119(3): 178–194.
- Boyla, K. & Estrada, A. Eds. (2005). Áreas Importantes para la Conservación de las Aves en los Andes Tropicales: Sitios prioritarios para la conservación de la biodiversidad. Quito, Ecuador: BirdLife International and Conservation International (BirdLife Conservation Series No. 14).
- Brown et al 2005
- Cisneros-Heredia, D.F. 2006. A preliminary approach to the Snipes (*Gallinago*) of Ecuador, with remarks on their distribution in Ecuadorian IBAs and its conservation status. *WI-WWSG Newsletter* 32: 4-11.
- Clay, R.P., A.J. Lesterhuis, S. Schulte, S. Brown, D. Reynolds & T.R. Simons. 2009. Conservation Plan for the American Oystercatcher (*Haematopus palliatus*) throughout the Western Hemisphere.
- Clay, R.P., Lesterhuis, A.J., Schulte, S., Brown, S., Reynolds, D. & Simons, T.R. 2014. A global assessment of the conservation status of the American Oystercatcher *Haematopus palliatus*. *International Wader Studies* 20: 62–82.
- Lesterhuis, A.J. & Clay, R.P. 2012. Synthesis of South American National Waterbird Reports. Unpub. Report to Waterbird Conservation Council.
- Lyons, J. E., W.L. Kendall, J.A. Royle, S.J. Converse, B.A. Andres and J.B. Buchanan. 2015. Population Size and Stopover Duration Estimation Using Mark–Resight Data and Bayesian Analysis of a Superpopulation Model. *Biometric* 72(1): 262–271.
- Wetlands International. 2013. "Waterbird Population Estimates 5". Retrieved from [wpe.wetlands.org](http://wpe.wetlands.org)
- Zdravkovic, M.G. 2013. Conservation Plan for the Wilson's Plover (*Charadrius wilsonia*). Version 1.0. Manomet Center for Conservation Sciences, Manomet, Massachusetts, USA.

**Compilado por:** Arne Lesterhuis, Oficina Ejecutiva de WHSRN